

# EFRON'S COINS AND THE LINIAL ARRANGEMENT

GÁBOR HETYEI

*To (Richard P. S. )<sup>2</sup>*

ABSTRACT. We characterize the tournaments that are dominance graphs of sets of (unfair) coins in which each coin displays its larger side with greater probability. The class of these tournaments coincides with the class of tournaments whose vertices can be numbered in a way that makes them semiacyclic, as defined by Postnikov and Stanley. We provide an example of a tournament on nine vertices that can not be made semiacyclic, yet it may be represented as a dominance graph of coins, if we also allow coins that display their smaller side with greater probability. We conclude with an example of a tournament with 81 vertices that is not the dominance graph of any system of coins.

## INTRODUCTION

A fascinating paradox in probability theory is due to B. Efron, who devised a four element set of nontransitive dice [3]: the first has three faces labeled 4 and two faces labeled 0, the second has all faces labeled 3, the third has four faces labeled 2 and two faces labeled 6, the fourth has three faces labeled 1 and three faces labeled 5. On this list, die number  $i$  defeats die number  $i + 1$  in the cyclic order in the following sense: when we roll the pair of dice simultaneously, die number  $i$  is more likely to display the larger number than die number  $i + 1$ . The paradox arose the interest of Warren Buffet, it inspired several other similar constructions and papers in game theory and probability: sample references include [1, 4, 5, 7].

This paper intends to investigate a hitherto unexplored aspect of Efron's original example: on each of its dice, only at most two numbers appear. They could be replaced with unfair coins, which display one of two numbers with a given probability. This restriction seems to be strong enough that we should be able to describe exactly which tournaments can be realized as *dominance graphs* of collections of unfair coins, where the direction of the arrows indicates which coin of a given pair is more likely to display the larger number.

Our paper gives a complete characterization in the case when we restrict ourself to the use of *winner* coins: these are coins that are more likely to display their larger number. The answer, stated in Theorem 3.2 is that a tournament has such a representation exactly when its vertices may be numbered in a way that it becomes a *semiacyclic tournament*. These tournaments were introduced by Postnikov and Stanley [9], the number of semiacyclic tournaments or  $n$  numbered vertices is the same as the number of regions of the  $(n - 1)$ -dimensional Linial arrangement. Our result allows to construct

---

*Date:* November 13, 2015.

*2010 Mathematics Subject Classification.* Primary 05C69; Secondary 60C5, 52C35, 05A.

*Key words and phrases.* nontransitive dice, semiacyclic tournament.

an example of a tournament on 9-vertices that can not be represented using winner coins only. On the other hand, we will see that this example is representable, if we also allow the use of *loser* coins, that is, coins that are more likely to display their smaller number. However, as we will see in Theorem 4.3, any tournament that is not representable with a set of winner coins only, gives rise via a direct product operation to a tournament that is not representable by any set of coins. In particular, we obtain an example of a tournament on 81 vertices that can not be represented by any set of coins as a dominance graph. Our results motivate several open questions, listed in the concluding Section 5.

## 1. PRELIMINARIES

**1.1. Hyperplane arrangements.** A hyperplane arrangement is a finite collection of codimension one hyperplanes in a finite dimensional vectorspace, together with the induced partition of the space into regions. The number of these regions may be expressed in terms of the Möbius function in the intersection lattice of the hyperplanes, using Zaslavsky's formula [10].

The *Linial arrangement*  $\mathcal{L}_{n-1}$  is the hyperplane arrangement

$$x_i - x_j = 1 \quad 1 \leq i < j \leq n \quad (1.1)$$

in the  $(n - 1)$ -dimensional vector space  $V_{n-1} = \{(x_1, \dots, x_n) \in \mathbb{R}^n : x_1 + \dots + x_n = 0\}$ . We will use the combinatorial interpretation of the regions of  $\mathcal{L}_{n-1}$  in terms of *semicyclic tournaments*, due to Postnikov and Stanley [9]. A *tournament* on the vertex set  $\{1, \dots, n\}$  is a directed graph with no loops nor multiple edges, such that for each 2-element subset  $\{i, j\}$  of  $\{1, \dots, n\}$ , exactly one of the directed edges  $i \rightarrow j$  and  $j \rightarrow i$  belongs to the graph. We may think of a tournament as the visual representation of the outcomes of all games in a championship, such that each team plays against each other team exactly once, and there is no “draw”.

**Definition 1.1.** A directed edge  $i \rightarrow j$  is called an ascent if  $i < j$  and it is a descent if  $i > j$ . For any directed cycle  $C = (c_1, \dots, c_m)$  we denote the number of directed edges that are ascents in the cycle by  $\text{asc}(C)$ , and the number of directed edges by  $\text{desc}(C)$ . A cycle is ascending if it satisfies  $\text{asc}(C) \geq \text{desc}(C)$ . A tournament on  $\{1, \dots, n\}$  is semicyclic if it contains no ascending cycle.

To each region  $R$  in  $\mathcal{L}_{n-1}$  we may associate a tournament on  $\{1, \dots, n\}$  as follows: for each  $i < j$  we set  $i \rightarrow j$  if  $x_i > x_j + 1$  and we set  $j \rightarrow i$  if  $x_i < x_j + 1$ . Postnikov and Stanley, and independently Shmulik Ravid, gave the following characterization of the tournaments arising this way [9, Proposition 8.5]

**Proposition 1.2.** A tournament  $T$  on  $\{1, \dots, n\}$  corresponds to a region  $R$  in  $\mathcal{L}_{n-1}$  if and only if  $T$  is semicyclic. Hence the number  $r(\mathcal{L}_{n-1})$  of regions of  $\mathcal{L}_{n-1}$  is the number of semicyclic tournaments on  $\{1, \dots, n\}$ .

## 2. THE COIN MODEL AND ITS ELEMENTARY PROPERTIES

In this paper we will study  $n$  element sets of (fair and unfair) coins. Each coin is described by a triplet of real parameters  $(a_i, b_i, x_i)$  where  $a_i \leq b_i$  and  $x_i > 0$  hold (here  $i = 1, 2, \dots, n$ ). The  $i$ th coin has number the number  $a_i$  on one side and  $b_i$  on the

other. After flipping it, it shows the number  $a_i$  with probability  $1/(1+x_i)$ , equivalently it shows the number  $b_i$  with probability  $x_i/(1+x_i)$ . Note that, as  $x_i$  ranges over the set of all positive real numbers, the probability  $1/(1+x_i)$  ranges over all numbers the open interval  $(0, 1)$ . We call the triplet  $(a_i, b_i, x_i)$  the *type* of the coin. We say that coin  $i$  *dominates* coin  $j$  if, after tossing both at the same time, the probability that coin  $i$  displays a strictly larger number than coin  $j$  is greater than the probability that coin  $i$  displays a strictly larger number. In other words, when we flip both coins, the one displaying the larger number “wins”, the other one “loses”, and we consider both coins displaying the same number a “draw”. The coin that is more likely to win, dominates the other.

We represent the domination relation as the *dominance graph*, whose vertices are the coins and there is a directed edge  $i \rightarrow j$  exactly when coin  $i$  dominates coin  $j$ . We will be interested in the question, which tournaments may be represented as the dominance graph of a set of  $n$  coins.

Up to this point we made one, inessential simplification: we assume that  $x_i$  can not be zero or infinity, that is, no coin can land on the same side with probability 1. If we have such a coin, we may replace it with a coin of type  $(a, a, 1)$ , that is, a fair coin that has the same number written on both sides. Since we are interested only in dominance graphs as tournaments we will also require that *for each pair of coins one dominates the other*. After fixing the parameters  $a_i$  and  $b_i$  for each coin, this restriction will exclude the points of  $\binom{n}{2}$  hypersurfaces of codimension one from the set of possible values of  $(x_1, x_2, \dots, x_n)$ , each hypersurface being defined by an equation involving a pair of variables  $\{x_i, x_j\}$ . The equations defining these surfaces may be obtained by replacing the inequality symbols with equal signs in the inequalities stated in Table 1 below. In particular, we assume that *different coins have different types*. In the rest of this section we will describe in terms of the types when the  $i$ th coin dominates the  $j$ th coin. To reduce the number of cases to be considered we first show that we may assume that no coin has the same number written on both sides. This is a direct consequence of the next lemma.

**Lemma 2.1.**

Suppose there is a coin of type  $(a_i, b_i, x_i)$ , satisfying  $a_i = b_i$ . Replace this coin with a coin of type  $(a'_i, b_i, 1)$  where  $a'_i$  is any real number that is less than  $a_i$  but larger than any element in the intersection of the set  $\{a_1, b_2, a_2, b_2, \dots, a_n, b_n\}$  with the open interval  $(-\infty, a_i)$ . Then modified system of coins has the same dominance graph.

*Proof.* We only need to verify that another coin  $j$ , of type  $(a_j, b_j, x_j)$  dominates a coin of type  $(a_i, a_i, x_i)$  if and only if it dominates a coin of type  $(a'_i, a_i, 1)$ . This is certainly the case when neither of  $a_j$  and  $b_j$  is equal to  $a_i$ , as these numbers compare to  $a_i$  the same way as to  $a'_i$ . We are left to consider the case when exactly one of  $a_j$  and  $b_j$  equals  $a_i$  (both can not equal because then there is no directed edge between  $i$  and  $j$  in the dominance graph).

**Case 1:**  $a_j = a_i$  and so  $a'_i < a_i = a_j < b_j$  hold. In the original system, as well as in the modified one, only coin  $j$  can win against coin  $i$  and it does so with a positive probability. We have  $j \rightarrow i$  in both dominance graphs.

**Case 2:**  $b_j = a_i$  and so  $a_j < a'_i < a_i = b_j$ . In the original system only coin  $j$  can lose (when it displays  $a_j$ ) so we have  $i \rightarrow j$  in the dominance graph. In the modified system,  $i$  can also lose sometimes, exactly when it displays  $a'_i$  (with probability  $1/2$ ) and coin  $j$  displays  $a_j$ . The probability of this event is  $1/(1+x_j)/2$ . A draw can also occur, exactly when both coins display  $a_i = b_j$ , the probability of this event is  $x_j/(1+x_j)/2$ . By subtracting these probabilities from 1 we obtain that the probability that  $j$  loses is exactly  $1/2$ . This is more than the probability of  $i$  losing, as  $1/(1+x_j)/2 < 1/2$ . Thus  $i \rightarrow j$  still holds in the dominance graph of the modified system.  $\square$

From now on we will assume that the type  $(a_i, b_i, x_i)$  of each coin satisfies  $a_i < b_i$ . If the type also satisfies  $x_i > 1$  then we call the coin a *winner*, if it satisfies  $x_i < 1$ , we call it a *loser*. Note that a loser coin can dominate a winner coin under the appropriate circumstances, these terms refer to the fact whether the coin is more likely to display its larger or smaller value. Note that a coin satisfies  $x_i = 1$  exactly when it is a *fair coin*.

In several results we will list our coins in *increasing lexicographic order of their types*.

**Definition 2.2.** We say that the type  $(a_i, b_i, x_i)$  is *lexicographically smaller* than the type  $(a_j, b_j, x_j)$ , if one of the following holds:

- (1)  $a_i < a_j$ ;
- (2)  $a_i = a_j$  and  $b_i < b_j$ ;
- (3)  $a_i = a_j$ ,  $b_i = b_j$  and  $x_i < x_j$ .

**Theorem 2.3.** Assume we list the coins in increasing lexicographic order by their types, comparing the coordinates left to right. Then, for  $i < j$ , we have  $i \rightarrow j$  if and only if exactly one of the following conditions is satisfied:

- (1)  $a_i = a_j < b_i < b_j$  and  $1/x_j > 1/x_i + 1$ ;
- (2)  $a_i < a_j < b_j < b_i$  and  $x_i > 1$ ;
- (3)  $a_i < a_j < b_i = b_j$  and  $x_i > x_j + 1$ ;
- (4)  $a_i < a_j < b_i < b_j$  and  $(1/x_i + 1)(x_j + 1) < 2$ .

*Proof.* Assuming that  $(a_i, b_i) \leq (a_j, b_j)$  holds in the lexicographic order, the six cases corresponding to the six lines of Table 1 below are a complete and pairwise mutually exclusive list of possibilities. The statement on the first line of Table 1 is obvious, and

Relation between $(a_i, b_i)$ and $(a_j, b_j)$	$i \rightarrow j$ exactly when
$(a_i, b_i) = (a_j, b_j)$	$x_i > x_j$
$a_i = a_j < b_i < b_j$	$1/x_j > 1/x_i + 1$
$a_i < a_j < b_j < b_i$	$x_i > 1$
$a_i < a_j < b_i = b_j$	$x_i > x_j + 1$
$a_i < a_j < b_i < b_j$	$(1/x_i + 1)(x_j + 1) < 2$
$a_i < b_i \leq a_j < b_j$	never

TABLE 1. Characterization of the dominance relation in terms of the types

$x_i > x_j$  can not happen as  $(a_i, b_i, x_i)$  comes before  $(a_i, b_i, x_j)$  in the lexicographic order exactly when  $x_i \leq x_j$  holds. The description on the third line is also clear, coin  $i$  wins exactly when it displays  $b_i$ , otherwise it loses. The last line is also obvious: coin  $j$  can

not lose in that case and it wins with a positive probability. The remaining lines require just a little more attention.

To prove the statement on the second line, observe first that coin  $i$  wins exactly when it displays  $b_i$  and coin  $j$  displays  $a_j$ , and it loses exactly when coin  $j$  displays  $b_j$  (regardless the outcome of tossing coin  $i$ ). Thus  $i \rightarrow j$  exactly when

$$\frac{x_i}{x_i + 1} \cdot \frac{1}{x_j + 1} > \frac{x_j}{x_j + 1}$$

which is equivalent to  $1/x_i + 1 < 1/x_j$ . The statement on the fourth line is completely analogous, only easier.

We are left to prove the statement on the fifth line. In this case, coin  $i$  wins exactly when it displays  $b_i$  and coin  $j$  displays  $a_j$ , and it loses in all other cases (there is never a draw). Therefore coin  $i$  dominates coin  $j$  if and only the probability of  $i$  winning is more than  $1/2$ , that is, we have

$$\frac{x_i}{x_i + 1} \cdot \frac{1}{x_j + 1} > \frac{1}{2}.$$

This is obviously equivalent to the statement on the last line. □

### 3. WINNER COINS REPRESENT SEMIACYCLIC TOURNAMENTS

In this section we give a complete description of all tournaments that may be realized as the domination graph of a system of coins that has only winner and fair coins. In proving our main result, the following lemma plays an important role.

**Lemma 3.1.** *Assume that the  $i$ th coin, of type  $(a_i, b_i, x_i)$  dominates the  $j$ th coin, of type  $(a_j, b_j, x_j)$  and that the  $j$ th coin is not a loser coin. Then we must have  $b_i \geq b_j$ .*

*Proof.* Assume by way of contradiction that  $b_j > b_i$  holds. The  $j$ th coin wins whenever it displays  $b_j$ , and this happens with probability  $x_j/(1 + x_j) \geq 1/2$ . We obtain that  $i \rightarrow j$  can not hold, a contradiction. □

**Theorem 3.2.** *Assume a set of  $n$  winner and fair coins is listed in increasing lexicographic order of their types. If the domination graph is a tournament, it must be semiacyclic. Conversely every semiacyclic tournament is the domination graph of a set of winner coins.*

*Proof.* Assume we are given an  $n$ -element set of winner and fair coins whose domination graph is a tournament and that the coins are listed in increasing lexicographic order of their types. Consider a cycle  $(i_1, i_2, \dots, i_k)$  in the domination graph. Repeated use of Lemma 3.1 yields  $b_{i_1} \geq b_{i_2} \geq \dots \geq b_{i_k} \geq b_{i_1}$ , implying  $b_{i_1} = \dots = b_{i_k}$ . As a consequence, using the fourth line in Table 1, we have  $x_{i_{j+1}} < x_{i_j} - 1$  whenever  $i_j < i_{j+1}$  and  $x_{i_{j+1}} < x_{i_j} + 1$  whenever  $i_{j+1} < i_j$ , for  $j = 1, \dots, k - 1$ . Similarly, we have  $x_{i_1} < x_{i_k} - 1$  whenever  $i_k < i_1$  and  $x_{i_1} < x_{i_k} + 1$  whenever  $i_1 < i_k$ . In other words, the last coordinate of the type increases by more than 1 at each ascent, and it increases by less than 1 at each descent. Therefore, a cycle can not be an ascending cycle, and the tournament must be semiacyclic.

Conversely, consider a semiacyclic tournament  $T$  on  $\{1, \dots, n\}$ . By Proposition 1.2 this tournament corresponds to a region of the Linial arrangement  $\mathcal{L}_{n-1}$  in the following way. For each  $i < j$  we have  $i \rightarrow j$  if  $x_i > x_j + 1$  and we have  $j \rightarrow i$  if  $x_i < x_j + 1$ , where  $(x_1, \dots, x_n)$  is an arbitrary point in the region. Note that some of the coordinates must be negative or zero here, as we have  $x_1 + \dots + x_n = 0$ . Introducing  $r := \max\{|x_1| + 1, \dots, |x_n| + 1\}$  and  $x'_i = x_i + r$ , we obtain a vector  $(x'_1, \dots, x'_n)$  whose coordinates are all positive, and satisfy  $i \rightarrow j$  if  $x'_i > x'_j + 1$  and  $j \rightarrow i$  if  $x'_i < x'_j + 1$ , for each  $i < j$ . Consider now the set of  $n$  coins where the  $i$ th coin has type  $(i, n+1, x'_i)$ . By the fourth line of Table 1, the dominance graph of this set of coins is precisely  $T$ .  $\square$

The proof of Theorem 3.2 also has the following consequence.

**Corollary 3.3.** *If a set of coins does not contain loser coins and its dominance graph is a tournament, then this tournament is also the dominance graph of a set of winner coins that all have the same number on one of their sides.*

Analogous results may also be stated for loser and fair coins. In analogy to Lemma 3.1 we can make the following observation.

**Lemma 3.4.** *Assume that the  $i$ th coin, of type  $(a_i, b_i, x_i)$  dominates the  $j$ th coin, of type  $(a_j, b_j, x_j)$  and that the  $i$ th coin is not a winner coin. Then we must have  $a_i \geq a_j$ .*

The proof is completely analogous and omitted. Lemma 3.4 may be used to prove the following result.

**Theorem 3.5.** *Assume a set of  $n$  loser and fair coins is listed in increasing lexicographic order of their types. If the domination graph is a tournament, it must be semiacyclic. Conversely every semiacyclic tournament is the domination graph of a set of loser coins.*

*Proof.* The proof is analogous to that of Theorem 3.2. Consider first the dominance graph of a set of  $n$  loser and fair coins and assume it is a tournament. We may use Lemma 3.4 to show that, in this dominance graph, all vertices contained in a cycle must have the same first coordinate. The role played by the fourth line of Table 1 is taken over by the second line, which may be used to show that the reciprocal of the third coordinate of the type increases by more than 1 after each ascent, and it decreases by less than 1 after each descent. Again we obtain that the dominance graph can not contain an ascending cycle.

Conversely, given a semiacyclic tournament  $T$  on the vertex set  $\{1, \dots, n\}$ , consider a point  $(x_1, \dots, x_n)$  satisfying  $x_1 + \dots + x_n = 0$ , such that for each  $i < j$  we have  $i \rightarrow j$  if  $x_i > x_j + 1$  and we have  $j \rightarrow i$  if  $x_i < x_j + 1$ . Let us set  $r := \max\{|1/x_1| + 1, \dots, |1/x_n| + 1\}$  and let us define  $x'_i$  by  $x'_i = r - 1/x_i$  for  $i = 1, \dots, n$ . Consider the set of  $n$  coins where the  $i$ th coin has type  $(0, i, x'_i)$  for  $i = 1, \dots, n$ . The dominance graph of this set of coins is  $T$ .  $\square$

In analogy to Corollary 3.3, the proof of Theorem 3.5 also has the following consequence.

**Corollary 3.6.** *If a set of coins does not contain winner coins and its dominance graph is a tournament, then this tournament is also the dominance graph of a set of loser coins that all have the same number on one of their sides.*

We conclude this section by an example of a tournament  $T$  whose vertices can not be labeled in an order that would make  $T$  semiacyclic. As a consequence,  $T$  can not be the dominance graph of a set of coins that does not contain winner, as well as loser coins. Our example will use a direct product construction, which we will reuse later, thus we make a separate definition.

**Definition 3.7.** *Given two tournaments  $T_1$  and  $T_2$  on the vertex sets  $V_1$  and  $V_2$  respectively, we define their direct product  $T_1 \times T_2$  as follows. Its vertex set is  $V_1 \times V_2$  and we set  $(u_1, v_1) \rightarrow (u_2, v_2)$  if either  $u_1 \rightarrow v_1$  belongs to  $T_1$  or we have  $u_1 = v_1$  and  $u_2 \rightarrow v_2$  belongs to  $T_2$ .*

This direct product operation is not commutative, but it is associative in the following sense: given  $k$  tournaments  $T_1, \dots, T_k$  on their respective vertex sets  $V_1, \dots, V_k$ , the vertex set of  $T_1 \times \dots \times T_k$  (where parentheses may be inserted in any order) is identifiable with the set of all  $k$ -tuples  $(v_1, \dots, v_k)$ , where, for each  $i$ ,  $v_i$  belongs to  $V_i$ . The edge  $(u_1, \dots, u_k) \rightarrow (v_1, \dots, v_k)$  belongs to  $T_1 \times \dots \times T_k$  exactly when  $u_i \rightarrow v_i$  belongs to  $T_i$  for the least  $i$  such that  $u_i \neq v_i$ .

**Proposition 3.8.** *Let  $C_3$  denote the 3-cycle. Then for any numbering of the vertex set of  $C_3 \times C_3$ , the resulting labeled tournament is not semiacyclic.*

*Proof.* Let us identify the vertices of  $C_3$  with 0, 1 and 2 in the order that  $0 \rightarrow 1, 1 \rightarrow 2, 2 \rightarrow 0$  belong to  $C_3$ . The vertex set of  $C_3 \times C_3$  is then the set of ternary strings of length 2. Assume, by way of contradiction, that there is a labeling of these ternary strings that makes  $C_3 \times C_3$  a semiacyclic tournament.

For each  $u \in \{0, 1, 2\}$  the cyclic permutation  $(u0, u1, u2)$  is an automorphism of the tournament  $(C_3 \times C_3)$ . Hence, without loss of generality, we may assume that in the set  $\{u0, u1, u2\}$  the vertex  $u0$  has the least label. As a consequence, the edge  $u0 \rightarrow u1$  is an ascent for each  $u$ . The cycle  $(00, 01, 10, 11, 20, 21)$  has 6 edges, and contains at least 3 ascents, thus it is an ascending cycle, in contradiction of having a semiacyclic tournament.  $\square$

#### 4. GENERAL SETS OF COINS

We begin with showing that the tournament  $C_3 \times C_3$  introduced in Proposition 3.8 can be represented with a set of coins, if we allow both winner and loser coins.

*Example 4.1.* We represent the vertices of  $C_3 \times C_3$  with the set of coins given in Table 2.

vertex	00	01	02
coin $(a, b, x)$	$(3, 6, 1 + \delta)$	$(2, 6, 3/2)$	$(1, 6, 2 + 2\delta)$
vertex	20	21	22
coin $(a, b, x)$	$(5, 12, 1/(s + 1 + 2\varepsilon))$	$(5, 11, 1/(s + 1/2))$	$(5, 10, 1/(s + \varepsilon))$
vertex	10	11	12
coin $(a, b, x)$	$(4, 9, 1/(r + 1 + 2\varepsilon))$	$(4, 8, 1/(r + 1/2))$	$(4, 7, 1/(r + \varepsilon))$

TABLE 2. A coin-representation  $C_3 \times C_3$

Here  $r$ ,  $s$ ,  $\varepsilon$  and  $\delta$  are positive real numbers, whose values we determine below. Initially we only require  $r > 1$  and  $s > 1$  which makes all coins loser except for the ones representing  $(0, 0)$ ,  $(0, 1)$  and  $(0, 2)$ .

To assure  $00 \rightarrow 01 \rightarrow 02 \rightarrow 00$  we need

$$2 + 2\delta < 3/2 + 1, \quad 3/2 < 1 + \delta + 1, \quad \text{and} \quad 2 + 2\delta > 1 + \delta + 1.$$

All three are satisfied if and only if we have

$$0 < \delta < 1/4. \tag{4.1}$$

To assure  $20 \rightarrow 21 \rightarrow 22 \rightarrow 20$  we need

$$r + 1/2 < r + \varepsilon + 1, \quad r + 1 + 2\varepsilon < r + 1/2 + 1, \quad \text{and} \quad r + 1 + 2\varepsilon > r + \varepsilon + 1.$$

All three are satisfied if and only if we have

$$0 < \varepsilon < 1/4. \tag{4.2}$$

Note that condition (4.2) is independent of the value of  $r$ , and we may replace  $r$  with  $s$  in the above calculations. Therefore to assure  $10 \rightarrow 11 \rightarrow 12 \rightarrow 10$  we also need to make sure (4.2), and this is a sufficient condition.

Note next, that any coin  $(4, b_1, x_1)$  representing a vertex of the form  $2v_1$  is dominated by any coin  $(5, b_2, x_2)$  representing a vertex of the form  $1v_2$ . Indeed, with probability  $1/(1 + x_1) > 1/2$  the first coin displays 4 and loses.

Next we make sure that any coin  $(4, b_1, x_1)$  representing a vertex of the form  $2v_1$  dominates any coin  $(a_2, 6, x_2)$  representing a vertex of the form  $0v_2$ . We have  $a_2 < 4 < 6 < b_1$  and, using line 5 of Table 1 we get that

$$(1 + 1/x_2)(1 + x_1) > 2$$

needs to be satisfied. Substituting the least value of  $x_1$  and the largest value of  $x_2$  we get

$$\left(1 + \frac{1}{2 + 2\delta}\right) \left(1 + \frac{1}{r + 1 + 2\varepsilon}\right) > 2$$

This inequality is equivalent to

$$r < \frac{2}{1 + \delta} - 2\varepsilon. \tag{4.3}$$

Finally, we want to make sure that any coin  $(5, b_1, x_1)$  representing a vertex of the form  $2v_1$  is dominated by any coin  $(a_2, 6, x_2)$  representing a vertex of the form  $0v_2$ . We have  $a_2 < 5 < 6 < b_1$  and, using line 5 of Table 1 we get that

$$(1 + 1/x_2)(1 + x_1) < 2$$

needs to be satisfied. Substituting the largest value of  $x_1$  and the least value of  $x_2$  we get

$$\left(1 + \frac{1}{1 + \delta}\right) \left(1 + \frac{1}{s + \varepsilon}\right) < 2.$$

This inequality is equivalent to

$$s > \frac{2 + \delta}{\delta} - \varepsilon. \tag{4.4}$$

The conditions (4.1), (4.2), (4.3) and (4.4) may be simultaneously satisfied, by setting  $\delta = 0.1$ ,  $\varepsilon = 0.1$ ,  $r = 1.6$  and  $s = 22$ , for example.

Example 4.1 proves that among the dominance graphs of systems of coins there are some that can not be labeled to become semiacyclic tournaments. On the other hand Theorems 3.2 and 3.5 imply an important necessary condition for a tournament to be the dominance graph of a system of coins.

**Corollary 4.2.** *If a tournament  $T$  may be represented as the dominance graph of a system of coins, then its vertex set  $V$  may be written as a union  $V = V_1 \cup V_2$ , such that the full subgraphs induced by  $V_1$  and  $V_2$ , respectively, may be labeled to become semiacyclic tournaments.*

Indeed, given a system of coins whose dominance graph is  $T$ , we may choose  $V_1$  to be the set of winner and fair coins and  $V_2$  to be the set of loser and fair coins (we do not have to include the fair coins on both sets) and then apply Theorems 3.2 and 3.5.

We conclude this section with an example of a tournament that can not be the dominance graph of any system of coins. Our example is  $C_3 \times C_3 \times C_3 \times C_3$ . The fact this tournament is not the dominance graph of any system of coins, of a direct consequence of Proposition 3.8 and the next theorem.

**Theorem 4.3.** *Suppose the tournaments  $T_1$  and  $T_2$  have the property that they are not semiacyclic for any ordering of their vertex sets. Then the tournament  $T_1 \times T_2$  can not be the dominance graph of any system of coins.*

*Proof.* Let us denote the vertex set of  $T_1$  and  $T_2$ , respectively, by  $V_1$  and  $V_2$ , respectively. Assume, by way of contradiction, that  $T_1 \times T_2$  is the dominance graph of a system of coins. As noted in Corollary 4.2, we may then write  $V_1 \times V_2$  as a union  $V_1 \times V_2 = W_1 \cup W_2$  such that the restriction of  $T_1 \times T_2$  to either of  $W_1$  or  $W_2$  can be labeled to become a semiacyclic tournament. For a fixed  $v_1 \in V_1$ , the restriction of  $T_1 \times T_2$  to the set  $\{(v_1, v_2) : v_2 \in V_2\}$  is isomorphic to  $T_2$ . Indeed, the first coordinate is the same for all vertices in the set and identifying each  $(v_1, v_2)$  with  $v_2$  yields an isomorphism. Since  $T_2$  can not be ordered to be semiacyclic, we obtain that  $\{(v_1, v_2) : v_2 \in V_2\}$  can not be entirely contained in  $W_2$ . As a consequence, for each  $v_1 \in V_1$  we can pick an  $f(v_1) \in V_2$  such that  $(v_1, f(v_1))$  belongs to  $W_1$ . Consider now the restriction of  $T_1 \times T_2$  to the set  $\{(v_1, f(v_1)) : v_1 \in V_1\}$ . This restriction is isomorphic to  $T_1$ , an isomorphism is given by  $(v_1, f(v_1)) \mapsto v_1$ . Since  $T_1$  can not be labeled to become a semiacyclic tournament, neither can the restriction of  $T_1 \times T_2$  to the set  $W_1$  that properly contains  $\{(v_1, f(v_1)) : v_1 \in V_1\}$ . We obtained a contradiction.  $\square$

## 5. CONCLUDING REMARKS

Theorem 3.2 motivates the question, how to describe those tournaments that can not be made semiacyclic, no matter how we order their vertices. Postnikov and Stanley have given a list of five cycles [9, Theorem 8.6], one of which must appear if the tournament (with numbered vertices) is not semiacyclic. Our question is different here, as we are allowed to choose our own numbering on the vertices.

**Conjecture 5.1.** *There is a finite list of tournaments that are minimally not semiacyclic in the sense, that removing any of their vertices allows the numbering of the remaining vertices in a semiacyclic way.*

We wish to make the analogous conjecture for dominance graphs of systems of coins.

**Conjecture 5.2.** *There is a finite list of tournaments that are minimally not dominance graphs of systems of coins in the sense, that after removing any of their vertices we obtain the dominance graph of a system of coins.*

Proving Conjecture 5.2 may be helped by looking at generalizations of hyperplane arrangements to hypersurface arrangements, that allow adding equations of hypersurfaces of the form  $1/x_j = 1/x_i + 1$  and of the form  $(1/x_i + 1)(x_j + 1) = 2$ , see Table 1. Such a study would only allow finding an analogue of semiacyclic tournaments (with numbered vertices), the question would still be open which subgraphs would obstruct numbering the vertices in a way that allows them to be represented in the desired way.

The number of regions in the Linial arrangement is listed as sequence A007889 in the OEIS [6]. These numbers count alternating trees and binary search trees as well. Several ways are known to find these numbers, see [2, 8, 9] neither of which seems to be related to counting semiacyclic tournaments directly. In view of the role played by these tournaments, it may be interesting to find a way to count them directly.

#### ACKNOWLEDGMENTS

This work was partially supported by a grant from the Simons Foundation (#245153 to Gábor Heteyei).

#### REFERENCES

- [1] N. Alon, G. Brightwell, H. A. Kierstead, A. V. Kostochka, and P. Winkler, Dominating sets in  $k$ -majority tournaments, *J. Combin. Theory Ser. B* **96** (2006), 374–387.
- [2] C. A. Athanasiadis, Extended Linial hyperplane arrangements for root systems and a conjecture of Postnikov and Stanley, *J. Algebraic Combin.* **10** (1999), 207–225.
- [3] M. Gardner, Mathematical Games: The Paradox of the Nontransitive Dice and the Elusive Principle of Indifference, *Sci. Amer.* **223** (1970), 110–114.
- [4] S. V. Gervacio, Severino and H. Maehara, Partial order on a family of  $k$ -subsets of a linearly ordered set, *Discrete Math.* **306** (2006), no. 4, 413–419.
- [5] R. Honsberger, Some Surprises in Probability, Ch. 5 in *Mathematical Plums* (Ed. R. Honsberger), Washington, DC: Math. Assoc. Amer., pp. 94–97, 1979.
- [6] OEIS Foundation Inc. (2011), “The On-Line Encyclopedia of Integer Sequences,” published electronically at <http://oeis.org>.
- [7] R. P. Savage, The Paradox of Nontransitive Dice, *The American Mathematical Monthly*, **101** (1994), 429–436.
- [8] A. Postnikov, Intransitive trees, *J. Combin. Theory Ser. A* **79** (1997), 360–366.
- [9] A. Postnikov and R. P. Stanley, Deformations of Coxeter hyperplane arrangements, *J. Combin. Theory Ser. A* **91** (2000), 544–97.
- [10] T. Zaslavsky, Facing up to arrangements: face-count formulas for partitions of space by hyperplanes, *Mem. Amer. Math. Soc.* **1** (1975), issue 1, no. 154, vii+102 pp.

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNC-CHARLOTTE, CHARLOTTE NC 28223-0001. WWW: <http://www.math.uncc.edu/~ghetyei/>.